




ELSEVIER

Contents lists available at ScienceDirect

DNA Repair

journal homepage: www.elsevier.com/locate/dnarepair

Characterization of rare NEIL1 variants found in East Asian populations

Irina G. Minko^a, Vladimir L. Vartanian^a, Naoto N. Tozaki^a, Oskar K. Linde^a, Pawel Jaruga^b, Sanem Hosbas Coskun^b, Erdem Coskun^b, Chunfeng Qu^c, Huan He^c, Chungui Xu^c, Taoyang Chen^d, Qianqian Song^c, Yuchen Jiao^c, Michael P. Stone^e, Martin Egli^f, Miral Dizdaroğlu^b, Amanda K. McCullough^{a,g}, R. Stephen Lloyd^{a,g,h,*}

^a Oregon Institute of Occupational Health Sciences, Oregon Health & Science University, Portland, OR, 97239, United States

^b Biomolecular Measurement Division, National Institute of Standards and Technology, Gaithersburg, MD, 20899, United States

^c State Key Lab of Molecular Oncology, National Cancer Center/National Clinical Research Center for Cancer/Cancer Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, 100021, Beijing, China

^d Qidong Liver Cancer Institute & Qidong People's Hospital, Qidong, 226200, Jiangsu Province, China

^e Department of Chemistry, Vanderbilt University, Nashville, TN, 37235, United States

^f Department of Biochemistry, Vanderbilt University, Nashville, TN, 37232, United States

^g Department of Molecular and Medical Genetics, Oregon Health & Science University, Portland, OR, 97239, United States

^h Department of Physiology and Pharmacology, Oregon Health & Science University, Portland, OR, 97239, United States

ARTICLE INFO

Keywords:

Base excision repair
Formamidopyrimidines
Thymine glycol
Aflatoxin
Hepatocellular carcinoma

ABSTRACT

The combination of chronic dietary exposure to the fungal toxin, aflatoxin B₁ (AFB₁), and hepatitis B viral (HBV) infection is associated with an increased risk for early onset hepatocellular carcinomas (HCCs). An in-depth knowledge of the mechanisms driving carcinogenesis is critical for the identification of genetic risk factors affecting the susceptibility of individuals who are HBV infected and AFB₁ exposed. AFB₁-induced mutagenesis is characterized by G to T transversions. Hence, the DNA repair pathways that function on AFB₁-induced DNA adducts or base damage from HBV-induced inflammation are anticipated to have a strong role in limiting carcinogenesis. These pathways define the mutagenic burden in the target tissues and ultimately limit cellular progression to cancer. Murine data have demonstrated that NEIL1 in the DNA base excision repair pathway was significantly more important than nucleotide excision repair relative to elevated risk for induction of HCCs. These data suggest that deficiencies in NEIL1 could contribute to the initiation of HCCs in humans. To investigate this hypothesis, publicly-available data on variant alleles of NEIL1 were analyzed and compared with genome sequencing data from HCC tissues derived from individuals residing in Qidong County (China). Three variant alleles were identified and the corresponding A51V, P68H, and G245R enzymes were characterized for glycosylase activity on genomic DNA containing a spectrum of oxidatively-induced base damage and an oligodeoxynucleotide containing a site-specific AFB₁-formamidopyrimidine guanine adduct. Although the efficiency of the P68H variant was modestly decreased, the A51V and G245R variants showed nearly wild-type activities. Consistent with biochemical findings, molecular modeling of these variants demonstrated only slight local structural alterations. However, A51V was highly temperature sensitive suggesting that its biological activity would be greatly reduced. Overall, these studies have direct human health relevance pertaining to genetic risk factors and biochemical pathways previously not recognized as germane to induction of HCCs.

1. Introduction

There exist a number of barriers to understanding the fundamental

genetic drivers that modulate susceptibility to the development of early onset hepatocellular carcinomas (HCCs) in individuals who are not only chronically exposed to the food-borne fungal toxin, aflatoxin B₁ (AFB₁),

Abbreviations: HCC, hepatocellular carcinoma; AFB₁, aflatoxin B₁; HBV, hepatitis B virus/viral; XP, xeroderma pigmentosum; NER, nucleotide excision repair; BER, base excision repair; AFB₁-FapyGua, 8,9-dihydro-8-(2,6-diamino-4-oxo-3,4-dihydropyrimid-5-yl-formamido)-9-hydroxyafatoxin B₁; FapyGua, 2,6-diamino-4-hydroxy-5-formamidopyrimidine; FapyAde, 4,6-diamino-5-formamidopyrimidine; ThyGly, thymine glycol; TAMRA, carboxytetramethylrhodamine; BHQ2, Black Hole Quencher 2; GC-MS/MS, gas chromatography-tandem mass spectrometry; Fpg, *E. coli* formamidopyrimidine glycosylase; Nth, *E. coli* endonuclease III

* Corresponding author.

E-mail address: lloydst@ohsu.edu (R.S. Lloyd).

<https://doi.org/10.1016/j.dnarep.2019.05.001>

Received 4 April 2019; Received in revised form 30 April 2019; Accepted 1 May 2019

Available online 03 May 2019

1568-7864/ © 2019 Published by Elsevier B.V.

but also infected with hepatitis B virus (HBV) (reviewed in [1] and [2]). Many extrinsic factors are known to be well-correlated with individuals who develop early-onset HCCs, including AFB₁, HBV and other factors which promote chronic inflammation. However, these environmental and lifestyle conditions are so ubiquitously distributed among at-risk populations in portions of East Asia, sub-Saharan Africa, Mexico, and Central and South America, that they are insufficient to predict individual susceptibility to the development of HCCs. Although it was anticipated that variants or altered gene expression of bioactivation and detoxication pathways or deficiencies in DNA nucleotide excision repair (NER) might contribute to individual susceptibility to the major AFB₁ DNA adducts, i.e. AFB₁-N7-guanine (AFB₁-N7-Gua) and two diastereomers of the imidazole ring-opened 8,9-dihydro-8-(2,6-diamino-4-oxo-3,4-dihydropyrimid-5-yl-formamido)-9-hydroxyafatoxin B₁ (AFB₁-FapyGua) [3–8], these hypotheses have generally not been supported by genome analyses except for rare homozygous inactivating mutations in xeroderma pigmentosum (XP) complementation groups C and D (XPC and XPD) NER enzymes [9,10]. In addition, studies demonstrating that XP complementation group A (XPA) knockout mice only exhibited a 1.2 risk ratio for the development of HCCs following maximally tolerated AFB₁ exposures cast doubt on the exclusive role of NER in the correction of AFB₁-induced DNA adducts [11]. These results, in combination with a paucity of data correlating specific allelic variants with increased susceptibility, suggest that the current understanding of the drivers in AFB₁- and inflammation-associated HCCs is inadequate to account for early onset disease.

Germane to these issues, we have discovered an additional DNA repair pathway for the removal of AFB₁-induced DNA damage which is of critical importance in limiting HCC formation. Specifically, we have demonstrated that NEIL1, a base excision repair (BER) DNA glycosylase/abasic site lyase, plays a key role in repair of AFB₁-FapyGua and that mice deficient in this enzyme are significantly more susceptible to AFB₁-induced HCCs [7]. These data raise the possibility that allelic variants of *NEIL1* which compromise catalytic activity or intracellular stability may modulate individual susceptibility to HCC formation. Furthermore, NEIL1 is also one of the key DNA glycosylases responsible for the removal of several oxidatively-induced DNA base lesions which are produced as a result of chronic inflammation. The major NEIL1 substrates include 2,6-diamino-4-hydroxy-5-formamidopyrimidine (FapyGua), 4,6-diamino-5-formamidopyrimidine (FapyAde) and thymine glycol (ThyGly) (reviewed in (12)). Collectively, these data identified NEIL1 as an essential enzyme for genome maintenance under conditions of both AFB₁ exposure and chronic inflammation.

We had demonstrated that the G83D and C136R variants of NEIL1 (Fig. 1) were glycosylase inactive [13], and suggested that catalytically or otherwise compromised NEIL1 variants could add to overall DNA adduct burdens sustained by hepatocytes. Relevant to this conclusion, recent data demonstrated that expression of the G83D variant of NEIL1 resulted in an oncogenic phenotype in MCF10A immortal human breast epithelial cells [14]. Expression of this variant, in the presence of WT NEIL1, induced replication stress, genomic instability, and cellular transformation through a mechanism postulated to involve the masking of DNA adducts at replication forks that are substrates for NEIL1. These data suggest that repair of AFB₁- and oxidatively-induced DNA damage could be blocked by catalytically-compromised, but correctly folded, variants of NEIL1 that are capable of binding to specific DNA base lesions. Additionally, we have previously demonstrated a significant phenotype in *Neil1*^{+/-} mice that showed intermediate (relative to WT and knockout) manifestations of the metabolic syndrome, including obesity, fatty liver disease, and elevated circulating leptin levels [15]. These data have significant implications for human health by linking NEIL1 functional heterozygosity with increased susceptibility to HCCs caused by AFB₁ or oxidatively-induced DNA damage.

Given the sum of biochemical and rodent data demonstrating the ability of NEIL1 to efficiently remove oxidatively-induced DNA base lesions and chemically stable AFB₁-FapyGua adducts, and considering




Fig. 1. Location of amino acid residues in NEIL1 that are changed in the variant alleles. Structure of NEIL1 (truncated form lacking 95 amino acids at the C-terminal domain [20]) with the protein depicted as a ribbon cartoon and in rainbow coloring from blue, N-terminus, to red, C-terminus. The key catalytic residue (P2) and the relevant residues that are changed in the variant alleles (G83, C136, A51, P68, and G245) are highlighted in ball-and-stick mode and labeled.

its critical role in protecting against AFB₁-induced HCCs in mice, this investigation was designed to identify and characterize the major NEIL1 variants in East Asia.

2. Materials and methods

2.1. Expression constructs

The variant alleles coding for full-length NEIL1 enzymes were constructed using the Q5 Site-Directed Mutagenesis Kit (New England Biolabs). The primer oligodeoxynucleotides (Integrated DNA Technologies) had the following sequences, with the variant nucleotide sites underlined: GCTTCAGTCCGCGGCAAGGA (A51V forward), TGAGATGCGGTAGGCACTGCT (A51V reverse); CCCAGC^uCCAACAGGAGCC (P68H forward); CCCAGGCAGAGGGCTCAG (P68H reverse); GGCTCAGGTCAGAGACGGG (G245R forward); CCTGCC^uCCCACTGGA CCA (G245R reverse). All designed substitutions were confirmed by DNA sequence analysis of the open reading frame for NEIL1 with a C-terminal 6-His-tag (DNA Sequencing Core, Vollum Institute, Oregon Health & Science University).

2.2. Expression and purification of NEIL1 enzymes

The expression vectors described above were introduced into BL21(DE3) *Escherichia coli* (New England Biolabs) and NEIL1 enzymes were purified as previously described [13]. Specifically, an overnight culture was prepared from a single colony in lysogeny broth supplemented with 50 µg/mL ampicillin at 37 °C, the culture was diluted 100-fold and continued to grow under these conditions to an optical density (OD₆₀₀) of 0.6. Expression of NEIL1 was induced by addition of isopropyl 1-thio-β-D-galactopyranoside to 0.5 mmol/L at 20 °C for 3.5 h. Cells were harvested and resuspended in 50 mmol/L sodium phosphate, pH 7.5, 300 mmol/L sodium chloride, 50 mmol/L imidazole (binding buffer) with EDTA-free complete protease inhibitor cocktail (Roche). Single-cell suspensions were passed through a French pressure cell at 8300 kPa. Cellular debris was removed by centrifugation at 10,000g for 20 min and the supernatant loaded onto a Ni-NTA agarose column (Qiagen), pre-equilibrated with binding buffer. The matrix was washed with ≈ 20 times the column volume of binding buffer, and proteins were eluted with a 50 mmol/L to 500 mmol/L imidazole gradient in a

total volume of 12-times the column volume and 2–3 mL fractions collected. The NEIL1-containing fractions were combined and dialyzed against 20 mmol/L Tris–HCl (pH 7.4), 100 mmol/L KCl, 1 mmol/L β -mercaptoethanol, 0.1 mmol/L EDTA, 15% (v/v) glycerol.

2.3. Site-specifically modified DNA substrates

The double-stranded ThyGly-containing DNA substrates conjugated with the carboxytetramethylrhodamine (TAMRA) fluorescent reporter were prepared from commercially produced oligodeoxynucleotides (Integrated DNA Technologies) according to the previously described protocol [16]. The adducted 17-mer oligodeoxynucleotide (5'-TCACC-G \underline{X} CGTACGACTC-3', where \underline{X} shows the location of ThyGly) had the TAMRA moiety on its 5' terminus. In assays using a plate reader for monitoring DNA incisions, the complement oligodeoxynucleotide (5'-GAGTCGTACGACGGTGA-3') contained the Black Hole Quencher 2 (BHQ2) on its 3' terminus. In the gel-based analyses, the complement strand had no modifications.

The 24-mer oligodeoxynucleotide containing a site-specific AFB₁-FapyGua was synthesized as reported [17]. The sequence of modified DNA was 5'-ACCACTACTAT \underline{X} ATTCATAACAAC-3', where \underline{X} corresponds to the adducted site. The adducted oligodeoxynucleotides were labeled at the 5' termini using [γ -³²P]ATP (PerkinElmer Life Sciences) and T4 polynucleotide kinase (New England Biolabs) and annealed with the complement strand (5'-GTTGTTATGAATCATAGTAGTGGT-3') as described [7].

2.4. DNA Glycosylase assays using site-specifically modified DNA substrates

The NEIL1-catalyzed reactions were conducted in 20 mmol/L Tris–HCl, pH 7.4, 100 mmol/L KCl, 100 μ g/mL BSA, and 0.01% (v/v) Tween 20. The detailed description of the experimental approach using TAMRA-conjugated DNA substrates and detection of the TAMRA fluorescent signal by a plate reader to monitor for DNA incisions was previously reported [16]. Briefly, DNA substrate with the TAMRA moiety on the 5' terminus of ThyGly-containing strand and the BHQ2 on the 3' terminus of the complement strand was aliquoted into wells of a black 384-well plate, and reactions were initiated by the addition of NEIL1 enzymes. The volume of a standard reaction was 20 μ L, DNA and enzyme were present at equimolar concentrations (50 nmol/L). Plates were incubated at 37 °C in the Infinite M200 plate reader (Tecan Group Ltd.). The result of DNA incision with subsequent dissociation of the TAMRA-conjugated product from the BHQ2-containing strand was an increase in fluorescent signal that was detected using a 525/9 nm excitation filter and a 598/20 nm emission filter. The data were recorded in arbitrary fluorescence units. In a subset of these experiments, formation of the incision products was validated using gel-based assays described below. These analyses revealed that at equimolar DNA and WT NEIL1 concentrations, the fluorescence plateau values correspond to complete incision of the substrate DNAs. However, there was a consistent, but slight delay, in the build-up of the fluorescence signal relative to DNA incision. Thus, the results of assays using a plate reader are presented in arbitrary fluorescence units, as opposed to percentages of product formed.

In gel-based assays, products were resolved through a 15% polyacrylamide gel under denaturing conditions (gel containing 8 mol/L urea and DNA samples containing 63% (v/v) formamide). For TAMRA-conjugated DNA, gel images were captured with the FluorChem M system (Protein Simple) using a 534 nm LED light source and 593 nm emission filter. The intensities of DNA bands were measured using Image Studio™ Lite version 4.0 software (LI–COR). The ³²P-labeled DNA was visualized using a Personal Molecular Imager™ System (Bio-Rad) and quantified from a phosphor screen image by the Quantity One Software (Bio-Rad).

The rate constants for NEIL1-catalyzed incisions were measured under single turnover conditions as previously described [7]. Following

pre-incubation at 37 °C for 2 min, the ³²P-labeled AFB₁-FapyGua substrate and enzyme were combined, incubated at 37 °C, and the aliquots removed during the course of reaction. The standard reaction contained 20 nmol/L double-stranded DNA substrate, with enzyme being present at 20- or 25-fold excess molar concentration. The aliquots were removed in the course of reactions, and in order to analyze glycosylase activity independently of lyase activity, DNA was incubated in the presence of 250 mmol/L NaOH at 90 °C for 5 min. Following gel electrophoresis and determination of band intensities, the product formation (P) was plotted as a function of time (t), and rate constant (k_{obs}), non-specific product (P_{ns}), and extrapolated amplitude of substrate used (S) were obtained from the best fit of the data to equation $P = P_{\text{ns}} + S(1 - e^{-k_{\text{obs}}t})$ using KaleidaGraph software (Synergy Software). The average values with standard deviations were calculated from three independent experiments.

2.5. Stabilities of NEIL1 enzymes at 37 °C

The NEIL1 enzymes were diluted in the reaction buffer described above to 100 nmol/L and incubated at 37 °C for 20 min. Incubations were monitored at 5 min intervals, tubes were placed on ice, and aliquots were collected and tested for activity using TAMRA/BHQ2-conjugated ThyGly substrate at equimolar enzyme to DNA concentrations (50 nmol/L). The production of fluorescent signal was monitored in a plate reader for 20 min and the rates were calculated from the linear phase of reactions (first 5 min). To assess the initial glycosylase activity, similar reactions also were conducted prior to incubation of enzyme at 37 °C. The rates obtained from the latter reactions were used as a reference to calculate the fraction of activity that remained at the given time point (A(t)). The average A(t) values from at least three independent experiments were plotted as a function of time (t), and rates of reduction in activity (λ) were obtained from the best fit of the data to equation $A(t) = e^{-\lambda t}$; the half-life was calculated using $t_{1/2} = \ln(2)/\lambda$.

2.6. Preparation of high-molecular weight DNA substrate for glycosylase reactions

Commercially available high-molecular weight calf thymus DNA (Sigma-Aldrich, St. Louis, MO) was dissolved in phosphate buffer (pH 7.4) at a concentration of 0.3 mg/mL at 4 °C. DNA solution was saturated with N₂O for 1 h and then irradiated in a ⁶⁰Co γ -ray source at a dose of 5 Gy, followed by dialysis against water for 18 h at 4 °C. Water outside the dialysis tubes was replaced twice. Aliquots of 50 μ g DNA were dried in a Speedvac under vacuum and kept at 4 °C until use.

2.7. NEIL1-catalyzed reactions using high-molecular weight DNA substrate and detection of excised DNA base lesions by gas chromatography-tandem mass spectrometry (GC–MS/MS)

For NEIL1-catalyzed reactions, aliquots of γ -irradiated DNA (50 μ g) were supplemented with stable isotope-labeled internal standards, i.e., FapyAde-¹³C, ¹⁵N₂, FapyGua-¹³C, ¹⁵N₂, 8-OH-Ade-¹³C, ¹⁵N₂, 8-OH-Gua-¹⁵N₂, 5-OH-Cyt-¹³C, ¹⁵N₂, 5-OH-Ura-¹³C, ¹⁵N₂, ThyGly-²H₄, 5-OH-5-MeHyd-¹³C, ¹⁵N₂ and 5,6-diOH-Ura-¹³C, ¹⁵N₂ (isodialuric acid-¹³C, ¹⁵N₂) which are a part of the NIST Standard Reference Material 2396 Oxidative DNA Damage Mass Spectrometry Standards (for details see <http://www.nist.gov/srm/> and https://www-s.nist.gov/srmors/view_detail.cfm?srm=2396). The samples were then dissolved in 50 μ L of 50 mmol/L phosphate buffer (pH 7.4), 100 mmol/L KCl, 1 mmol/L EDTA and 0.1 mmol/L dithiothreitol. Samples were incubated for 1 h at 37 °C using enzyme concentrations that are indicated in the figure legend. In addition to reactions in the presence of NEIL1 enzymes, control reactions were performed in parallel that had no enzyme or contained a combination of the *E. coli* formamidopyrimidine glycosylase (Fpg) (R&D Systems, Inc. Minneapolis, MN) and endonuclease III (Nth) (Prospect-Tany TechnoGene Ltd., Ness Ziona, Israel). Following precipitation of

DNA with ethanol, the supernatant fractions were collected. After removal of ethanol in a Speedvac, the samples were lyophilized and trimethylsilylated. Specifically, 60 μ L of a mixture of nitrogen-bubbled bis(trimethylsilyl)trifluoroacetic acid [containing trimethylchlorosilane (1%; v/v)] (BSTFA) (Pierce, Rockford, Ill.) and pyridine (Sigma-Aldrich, St. Louis, MO) (1:1, v/v) was added to each lyophilized supernatant fraction. The samples were vortexed and purged individually with ultra-high-purity nitrogen, sealed with Teflon-coated septa, and heated at 120 °C for 30 min. After cooling, the clear samples were removed and placed in vials used for injection onto the GC-column. Aliquots (4 μ L) of derivatized samples were analyzed by gas chromatography-tandem mass spectrometry (GC-MS/MS) under the conditions described previously [18]. On the basis of the known mass spectra of trimethylsilyl derivatives of DNA base lesions (reviewed in [19]), the following mass transitions were used for identification and quantification: m/z 369 \rightarrow m/z 280 and m/z 372 \rightarrow m/z 283 for FapyAde and FapyAde- ^{13}C , $^{15}\text{N}_2$ respectively; m/z 457 \rightarrow m/z 368 and m/z 460 \rightarrow m/z 371 for FapyGua and FapyGua- ^{13}C , $^{15}\text{N}_2$, respectively; m/z 367 \rightarrow m/z 352 and m/z 370 \rightarrow m/z 355 for 8-OH-Ade and 8-OH-Ade- ^{13}C , $^{15}\text{N}_2$, respectively; m/z 455 \rightarrow m/z 440 and m/z 460 \rightarrow m/z 445 for 8-OH-Gua and 8-OH-Gua- $^{15}\text{N}_5$, respectively; m/z 343 \rightarrow m/z 342, and m/z 346 \rightarrow m/z 345 for 5-OH-Cyt and 5-OH-Cyt- ^{13}C , $^{15}\text{N}_2$, respectively; m/z 344 \rightarrow m/z 343, and m/z 350 \rightarrow m/z 349 for 5-OH-Ura and 5-OH-Ura- $^{13}\text{C}_4$, $^{15}\text{N}_2$, respectively; m/z 448 \rightarrow m/z 259 and m/z 452 \rightarrow m/z 262 for ThyGly and ThyGly- $^2\text{H}_4$, respectively. The quantification was achieved using integrated areas of the signals of the mass transitions of the monitored DNA base lesions and their corresponding stable isotope-labeled analogues. The measured levels were expressed as number of lesions per 10^6 DNA bases. The mean values with standard deviations were calculated from three independent measurements. Significant excision of FapyGua, FapyAde and ThyGly from DNA was observed, whereas the other DNA lesions measured in this work were not significantly excised by NEIL1 enzymes over control levels.

2.8. Modeling the structural consequences of NEIL1 A51V, P68H and G245R allelic variants

The crystallographic coordinates for C-terminally truncated NEIL1 (Δ 95-CTD) [20] were retrieved from the Protein Data Bank (www.rcsb.org; PDB ID 5ITQ). Amino acids A51, P68 and G245 were swapped to V, H, and R, respectively, in UCSF Chimera [21]. The unfavorable *cis* conformation of H68 following mutation from proline was converted to *trans* using the program Coot [22], followed by real-space refinement inside the confines of the final electron density for the NEIL1 crystal structure. The mutated protein structure was then energy minimized with Amber 14 [23] using steepest descent and conjugate gradient cycles until convergence.

3. Results and discussion

3.1. NEIL1 allelic variants in East Asian populations

Previous data from our laboratories have revealed that in addition to NER being important in the repair of AFB₁-induced DNA adducts, the BER pathway is of critical importance in limiting HCC formation by the removal of AFB₁-induced DNA damage (reviewed in (2)). These data raise the possibility that allelic variants of NEIL1 with compromised catalytic activity or intracellular stability may modulate individual susceptibility to AFB₁-induced HCC. Consequently, we have hypothesized that genetic variation in NEIL1 may modulate HCC susceptibility arising from both AFB₁ exposure and high frequency HBV infection.

To investigate this hypothesis, we have carried out systematic analyses of publicly-available DNA sequence data bases for NEIL1 variant allele frequencies specifically using the Genome Aggregation Database (gnomAD) [24] and the 1000 Genomes [25]. The gnomAD contains ~300,000 alleles representing six global and eight sub-

continental ancestries. The previously characterized glycosylase-incompetent NEIL1 variants were not found in East Asian populations; G83D is associated with European, Ashkenazi Jewish, and Latino ancestries and C136R is found in African populations and at extremely low frequencies. When singletons were excluded, a total of 20 missense variants in the exomes of NEIL1 were identified in East Asian populations, with 3 of these, A51V (rs768187848), P68H (rs187873972), and G245R (rs140982397), present at > 0.1% frequencies. The positions of these residues within the NEIL1 structure are shown in Fig. 1. Subsequent analyses of the 1000 Genomes database [25] revealed that the most common East-Asian variant, P68H, which presents in this region at an overall frequency of \approx 1.4%, is overrepresented in some ethnic groups. Specifically, eight and six alleles for this variant were found in exomes of 190 individuals from Vietnam (Kinh in Ho Chi Minh City) and 180 individuals from China (Chinese Dai in Xishuangbanna), respectively.

Our hypothesis may also be supported by exome sequencing data of HCC patients who reside in a region of Southeast China with known AFB₁ exposure [26]. In this data set, each of the three NEIL1 variant alleles (A51V, P68H, and G245R) was identified once in the 49-patient cohort. These findings are notable and unanticipated because individually, these allelic variants are only present in East Asia at \approx 0.17%, 1.4%, and 0.65%, respectively. Further population-based studies are necessary to test the hypothesis that there is an association between rare genetic variations in the NEIL1 gene and susceptibility to HCC.

3.2. Substrate specificity of WT and allelic variants of NEIL1

To express proteins encoded by NEIL1 nonsynonymous allelic variants (A51V, P68H, and G245R), the full-length coding sequence for each variant was engineered into an expression plasmid for K242R NEIL1. The K242R version of NEIL1 results from editing of pre-mRNA by adenosine deaminase, ADAR1 that converts the central adenosine in an AAA codon to inosine, thus altering the translation code from K to R [27]. The initial characterization of human NEIL1 [28–30], as well as the majority of subsequent studies on NEIL1 biochemical properties, structure, and modulation of its activity by amino acid changes has been done using the edited version. Thus, WT NEIL1 and its variants investigated here were represented by the edited, K242R forms. All plasmids were sequence verified prior to protein expression in *E. coli*, and each was purified using Ni-agarose chromatography. Protein preparations were determined to be 90% to 95% pure, with no non-specific nicking activities on control DNA.

Preliminary kinetic analyses were carried out on fluorescently-labelled synthetic oligodeoxynucleotides in which a 5' TAMRA-labelled 17-mer, containing a ThyGly base at position 7, was annealed with a complementary strand containing a 3' BHQ2 moiety [16]. Reactions were initiated by addition of WT or variant proteins to DNA substrate and analyzed in real-time in a plate reader. Since the site of DNA modification was 6 nucleotides from the 5' TAMRA label, incision of substrate resulted in the melting of the 6-mer and an enhancement of fluorescence. As shown in Fig. 2, all three variant enzymes could cleave the ThyGly-containing oligodeoxynucleotide. The rate of the fluorescent signal production in the presence of A51V (Fig. 2A) was comparable with that of WT during the early phase of reaction (\approx first 10 min), but significantly slowed soon thereafter. This kinetics contrasted with a parallel reaction with WT, in which the fluorescent signal continued to increase throughout the time course. The reaction kinetics observed for P68H (Fig. 2B) and G245R (Fig. 2C) were indistinguishable relative to that of WT. These results demonstrating the abilities of A51V, P68H, and G245R to cleave ThyGly-containing oligodeoxynucleotide were corroborated using gel-based assays (data not shown) and in relation to G245R, agreed with results of a previous study [31]. Thus, all three NEIL1 variants possess glycosylase activity for ThyGly and retain lyase function.




Fig. 2. Initial characterization of the NEIL1 variants using fluorescently-labeled ThyGly-containing oligodeoxynucleotides. The activities of the A51V (A), P68H (B), and G245R (C) variants were tested along with activity of WT NEIL1 using TAMRA/BHQ2-conjugated double-stranded oligodeoxynucleotides that contained a site-specific ThyGly. The reactions were conducted with equimolar DNA and enzyme concentrations (50 nmol/L) at 37 °C. The TAMRA emission was recorded using an Infinite M200 plate reader.




Fig. 3. Removal of oxidatively-induced DNA base lesions from high-molecular weight DNA by WT and variants of NEIL1. The γ -irradiated calf thymus DNA (50 μ g) was incubated for 1 h at 37 °C in the presence of NEIL1 enzymes (2 μ g), combination of Fpg and Nth (1 μ g each), or in reaction buffer. The released FapyGua (A), FapyAde (B), and ThyGly (C) were measured by GC–MS/MS using their stable isotope-labeled analogues as internal standards. The statistically significant differences are only shown for variant enzyme versus WT (* = $p < 0.05$; ** = $p < 0.01$). The differences between the non-specific base hydrolysis and the base release in the presence of enzymes were statistically significant in all cases ($p < 0.05$). The uncertainties are standard deviations.

Although the analyses of incision of ThyGly-containing synthetic oligodeoxynucleotides provided initial insights into the activity of these enzymes, the substrate specificity of NEIL1 is more broad and includes FapyGua and FapyAde as the primary base lesions excised from genomic DNAs containing a spectrum of oxidatively-induced DNA damages (reviewed in [12]). Thus, purified WT and each of the allelic variants of NEIL1 were incubated with calf thymus DNA that had been previously exposed to a 5 Gy γ -irradiation dose. Control reactions (no enzyme treatment) and treatment with a combination of the *E. coli* Fpg and Nth were also performed. Fpg has been previously characterized to release FapyGua, FapyAde and 8-hydroxyguanine with similar excision kinetics [32–34], while Nth has a broad substrate specificity for pyrimidine-derived lesions, including ThyGly, and also removes FapyAde [35,36]. Relative to the WT NEIL1, the efficiency of P68H was reduced on both FapyGua and FapyAde, while the efficiency of G245R was only reduced relative to the release of FapyGua (Fig. 3A and B). Consistent with the results of experiments with site-specifically modified oligodeoxynucleotides, P68H and G245R released ThyGly from genomic DNA with the same efficiency as WT NEIL1, while A51V showed a modest increase in the efficiency of ThyGly incision (Fig. 3C). These data on the excision of ThyGly from genomic DNAs emphasize that this damage is only poorly recognized by edited NEIL1 [37] relative to the total ThyGly in the sample as revealed by the combined Nth/Fpg treatment. The poor excision of ThyGly was in stark contrast to the

efficient excision of both FapyGua and FapyAde by NEIL1 enzymes; these base lesions were released at levels comparable to that observed in Fpg/Nth-catalyzed reactions (Fig. 3A and B).

3.3. Temperature sensitivity of the A51V NEIL1 variant

The kinetics of A51V-catalyzed reaction with ThyGly-containing oligodeoxynucleotides (Fig. 2A) suggested that incision was nearly complete ≈ 10 min into the reaction. One possible explanation is that the A51V allelic variant is temperature sensitive and that during the reaction, there was an appreciable loss in catalytic activity. To test this hypothesis, WT and each of the variants were incubated at 37 °C for increasing times in reaction buffer that did not contain the DNA substrate and the activities were evaluated from the linear phase of reaction following the addition of damage-containing oligodeoxynucleotides. The fraction of activity remaining relative to the initial activity was plotted as a function of pre-incubation time (Fig. 4) and the experimental points were fit to an exponential equation to obtain rates of reduction in activity. Consistent with the hypothesis of A51V being temperature sensitive, the half-life at 37 °C was estimated to be ≈ 5 min versus ≈ 30 min estimated for WT (Fig. 4A). The gel-based analyses confirmed a rapid inactivation of A51V at 37 °C (data not shown). The A51V enzyme that had been inactivated for 20 min at 37 °C was kept in ice for 1 h and 24 h, and its activity was retested. No evidence for the




Fig. 4. Kinetics of temperature inactivation of WT and variants of NEIL1. WT and variants of NEIL1 A51V (A), P68H (B), and G245R (C) were pre-incubated at 37 °C and the activities were tested with equimolar DNA and enzyme concentrations (50 nmol/L) using TAMRA/BHQ2-conjugated double-stranded ThyGly substrate. The reactions were conducted at 37 °C and monitored by an Infinite M200 plate reader. The rates were calculated from the linear phase of reactions. The fraction of activity remaining at the given time point ($A(t)$) was calculated relative to the initial enzyme activity. The average $A(t)$ values from at least three independent experiments were plotted as function of time (t). The uncertainties are standard deviations.

activity being restored was found (data not shown), suggesting that inactivation of this variant under physiological temperature conditions is irreversible. In contrast, P68H and G245R displayed no enhanced temperature inactivation relative to that of the WT enzyme (Fig. 4B and C, respectively).

Considering a poor recognition of ThyGly by the edited NEIL1 enzymes (Fig. 3C), an experiment was designed to validate thermal instability of A51V using the preferred NEIL1 substrates, FapyGua and FapyAde. The A51V variant was pre-incubated at 37 °C for 30 min, and then reacted with γ -irradiated calf thymus DNA. The levels of the released base lesions were assessed by GC–MS/MS (Fig. 5). The data confirmed enzyme inactivation under physiological temperature conditions. The removal of FapyGua and FapyAde was reduced 7-fold and 5-fold, respectively (Fig. 5A and B) and consistent with the results of experiments with synthetic oligodeoxynucleotides (Fig. 4A), no residual activity was detected with regard to excision of ThyGly (Fig. 5C).

3.4. Incision of synthetic oligodeoxynucleotides containing AFB₁-FapyGua

In addition to characterizing the activities of these allelic variants on DNAs containing oxidatively-induced base damages, we also characterized their efficiencies in the incision of DNA containing a site-specific AFB₁-FapyGua base adduct (Table 1 and Fig. 6). As shown in Table 1, P68H showed reduced rates of incision relative to WT, with the $p < 0.05$, while the rate of G245R-catalyzed reaction was comparable to that of WT. The analysis of A51V under single-turnover conditions was not practical due to incompatibility of its temperature sensitivity with conditions under which the single-turnover experiments were conducted. Thus, the ability of A51V to incise AFB₁-FapyGua was tested

Table 1

Single-turnover kinetics of WT and variants of NEIL1 on AFB₁-FapyGua-containing oligodeoxynucleotides. The uncertainties are standard deviations.

Enzyme	Incision rates (k_{obs} , min ⁻¹)
Wild type	0.18 ± 0.01
A51V	ND
P68H	0.13 ± 0.01
G245R	0.16 ± 0.02




Fig. 6. Incision of AFB₁-FapyGua from site-specifically modified oligodeoxynucleotides by WT and A51V NEIL1. The reactions were conducted with 500 nmol/L WT or A51V NEIL1 at 37 °C for 5 min using [³²P]-labeled 24-mer double-stranded DNA (20 nmol/L) that contained a centrally-located AFB₁-FapyGua. Representative gel image (A) and percent of product formation (average ± standard deviation) calculated from three independent experiments (B) are shown (* = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$). The uncertainties are standard deviations.




Fig. 5. Removal of oxidatively-induced DNA base lesions by A51V following pre-incubation of enzyme at 37 °C. The γ -irradiated calf thymus DNA (50 μ g) was incubated for 1 h at 37 °C in the presence of A51V enzyme (2 μ g). The aliquot of A51V was pre-incubated at 37 °C for 30 min. Subsequently, DNA was added to the solution which was then incubated for another 30 min. The released FapyGua (A), FapyAde (B), and ThyGly (C) were measured by GC–MS/MS using their stable isotope-labeled analogues as internal standards (* = $p < 0.05$; ** = $p < 0.01$; *** = $p < 0.001$). The uncertainties are standard deviations.




Fig. 7. NEIL1 crystal structure and conformational effects of allelic variants. Overall structure of NEIL1 (truncated form lacking 95 amino acids at the C-terminal domain [20]) with the protein depicted as a ribbon cartoon and in rainbow coloring from blue, N-terminus, to red, C-terminus, and variant amino acids highlighted in ball-and-stick mode and labeled (A). Overlay of the WT and variant structures viewed in the regions around A/V 51 (B), P/H 68 (C), and G/R 245 (D). Residues in the variant and energy-minimized structure are colored by atom, with carbon, oxygen and nitrogen atoms shown in tan, red and blue, respectively, and the WT structure is uniformly colored in gray. Variant residues 51, 68 and 245 are highlighted in ball-and-stick mode and selected neighboring residues are labeled.

in short, 5-min reactions and the incision product was observed (Fig. 6). However, given a short half-life of this variant at physiologically relevant temperature, the biological significance of this observation can be questioned.

3.5. Modeling the structural consequences of NEIL1 A51V, P68H and G245R allelic variants

The conformational changes as a result of amino acid substitutions are quite limited as per the computational model of A51V-P68H-G245R NEIL1 (Fig. 7). In the WT protein, amino acids P68 and G245 map to loop regions on the surface whereas A51 lies in the interior between a β -strand and an α -helix (Fig. 7A). In the latter case, only slight shifts are necessary to accommodate the bulkier valine side chain (Fig. 7B). The change from *cis*-amide proline to *trans*-amide histidine at position 68 only alters the conformation locally and main and side chain atoms in the WT and mutated NEIL1 are superimposable at the neighboring Q67 and Q70 (Fig. 7C). Likewise, the large arginine side chain in the loop at position 245 juts away from the protein surface and can be accommodated by slight adjustments in the preceding Y244 and following S246 and the neighboring residues F13 and Q86 (Fig. 7D).

The observed correlations between the locations of substituted amino acids, the structural changes as inferred from computational modeling, and the phenotypes of NEIL1 sequence variants follow general trends found in protein structure-function relationships. Thus, amino acid changes near the periphery are typically less deleterious than changes mapping to the active site or near the ligand and/or substrate [38]. In 3-methyladenine DNA glycosylase, mutations in surface loops, if not involved in DNA binding, are tolerated better relative to those buried in the protein's exterior, especially when present in β sheets [39]. Consistent with these prior observations, substitutions

in the surface loop regions, at P68 and G245, imposed only small local changes, with no or minor phenotypical consequences. Deviations from the WT structure in A51V were also minimal, but occurred at the interface of a β -strand and an α -helix. We hypothesize that these local changes are sufficient to facilitate enzyme misfolding, which is manifested in thermal instability. The structural basis for the severely affected phenotype of the G83D variant had previously been addressed. G83D was predicted to affect the conformation of the β 4/5 loop by introducing an unfavorable negative charge in an otherwise hydrophobic local environment [40]. Indeed, the displacement of residues of the β 4/5 loop was observed in the crystal structure of the corresponding G86D variant in Mimivirus Neil1 [37].

4. Conclusion

The goal of this investigation was to determine whether rare allelic variants of NEIL1 that are geographically clustered in East Asia display alterations in kinetics of excision, substrate specificities, and stabilities relative to the WT enzyme. Our data demonstrated that the P68H allelic variant had a modest reduction in catalytic activity on the release of FapyGua, FapyAde, and AFB₁-FapyGua. The substrate specificity and catalytic efficiency of G245R was indistinguishable from WT, except for a reduced efficiency in excision of FapyGua. Additionally, the A51V variant was unstable under physiological temperature conditions, suggesting that it may have no biological activity. Although the significance of these changes is yet to be determined in humans, the presence of these rare allelic variants in HCC patients from Qidong County is suggestive that there could be adverse health outcomes. These studies provide the framework from which larger, disease-based molecular epidemiology studies will be conducted.

Funding source

This study was supported in part by the National Institutes of Health (NIH) Grants R01 CA055678, R01 ES029357, and R56 ES027632. The work of Oskar K. Linde was partially supported by the Summer Internship Program (Oregon Institute of Occupational Health Sciences, Oregon Health & Science University).

Conflict of interest

None declared.

Transparency document

The [Transparency document](#) associated with this article can be found in the online version.

Acknowledgments

Certain commercial equipment or materials are identified in this paper in order to specify adequately the experimental procedure. Such identification does not imply recommendation or endorsement by the National Institute of Standards and Technology, nor does it imply that the materials or equipment identified are necessarily the best available for the purpose.

References

- [1] B.I. Fedeles, J.M. Essigmann, Impact of DNA lesion repair, replication and formation on the mutational spectra of environmental carcinogens: aflatoxin B₁ as a case study, *DNA Repair (Amst)* 71 (2018) 12–22.
- [2] A.K. McCullough, R.S. Lloyd, Mechanisms underlying aflatoxin-associated mutagenesis—implications in carcinogenesis, *DNA Repair (Amst)* 77 (2019) 76–86.
- [3] J.M. Essigmann, R.G. Croy, A.M. Nadzan, W.F. Busby Jr., V.N. Reinhold, G. Buchi, G.N. Wogan, Structural identification of the major DNA adduct formed by aflatoxin B₁ in vitro, *Proc. Natl. Acad. Sci. U. S. A.* 74 (1977) 1870–1874.

- [4] R.G. Croy, J.M. Essigmann, V.N. Reinhold, G.N. Wogan, Identification of the principal aflatoxin B₁-DNA adduct formed in vivo in rat liver, *Proc. Natl. Acad. Sci. U. S. A.* 75 (1978) 1745–1749.
- [5] R.G. Croy, G.N. Wogan, Temporal patterns of covalent DNA adducts in rat liver after single and multiple doses of aflatoxin B₁, *Cancer Res.* 41 (1981) 197–203.
- [6] L.L. Woo, P.A. Egner, C.L. Belanger, R. Wattanawaraporn, L.J. Trudel, R.G. Croy, J.D. Groopman, J.M. Essigmann, G.N. Wogan, J.T. Bouhenguel, Aflatoxin B₁-DNA adduct formation and mutagenicity in livers of neonatal male and female B6C3F1 mice, *Toxicol. Sci.* 122 (2011) 38–44.
- [7] V. Vartanian, I.G. Minko, S. Chawanthayatham, P.A. Egner, Y.C. Lin, L.F. Earley, R. Makar, J.R. Eng, M.T. Camp, L. Li, M.P. Stone, M.R. Lasarev, J.D. Groopman, R.G. Croy, J.M. Essigmann, A.K. McCullough, R.S. Lloyd, NEIL1 protects against aflatoxin-induced hepatocellular carcinoma in mice, *Proc. Natl. Acad. Sci. U. S. A.* 114 (2017) 4207–4212.
- [8] E. Coskun, P. Jaruga, V. Vartanian, O. Erdem, P.A. Egner, J.D. Groopman, R.S. Lloyd, M. Dizdaroglu, Aflatoxin-guanine DNA adducts and oxidatively-induced DNA damage in aflatoxin-treated mice in vivo as measured by liquid chromatography-tandem mass spectrometry with isotope-dilution, *Chem. Res. Toxicol.* 32 (2019) 80–89.
- [9] X.D. Long, Y. Ma, Y.F. Zhou, A.M. Ma, G.H. Fu, Polymorphism in xeroderma pigmentosum complementation group C codon 939 and aflatoxin B₁-related hepatocellular carcinoma in the Guangxi population, *Hepatology* 52 (2010) 1301–1309.
- [10] X.D. Long, Y. Ma, Y.F. Zhou, J.G. Yao, F.Z. Ban, Y.Z. Huang, B.C. Huang, XPD codon 312 and 751 polymorphisms, and AFB₁ exposure, and hepatocellular carcinoma risk, *BMC Cancer* 9 (2009) 400.
- [11] Y. Takahashi, Y. Nakatsuru, S. Zhang, Y. Shimizu, H. Kume, K. Tanaka, F. Ide, T. Ishikawa, Enhanced spontaneous and aflatoxin-induced liver tumorigenesis in xeroderma pigmentosum group A gene-deficient mice, *Carcinogenesis* 23 (2002) 627–633.
- [12] M. Dizdaroglu, E. Coskun, P. Jaruga, Repair of oxidatively induced DNA damage by DNA glycosylases: mechanisms of action, substrate specificities and excision kinetics, *Mutat. Res.* 771 (2017) 99–127.
- [13] L.M. Roy, P. Jaruga, T.G. Wood, A.K. McCullough, M. Dizdaroglu, R.S. Lloyd, Human polymorphic variants of the NEIL1 DNA glycosylase, *J. Biol. Chem.* 282 (2007) 15790–15798.
- [14] H.A. Galick, C.G. Marsden, S. Kathe, J.A. Dragon, L. Volk, A.A. Nemec, S.S. Wallace, A. Prakash, S. Double, J.B. Sweasy, The NEIL1 G83D germline DNA glycosylase variant induces genomic instability and cellular transformation, *Oncotarget* 8 (2017) 85883–85895.
- [15] V. Vartanian, B. Lowell, I.G. Minko, T.G. Wood, J.D. Ceci, S. George, S.W. Ballinger, C.L. Corless, A.K. McCullough, R.S. Lloyd, The metabolic syndrome resulting from a knockout of the NEIL1 DNA glycosylase, *Proc. Natl. Acad. Sci. U. S. A.* 103 (2006) 1864–1869.
- [16] A.C. Jacobs, M.J. Calkins, A. Jadhav, D. Dorjsuren, D. Maloney, A. Simeonov, P. Jaruga, M. Dizdaroglu, A.K. McCullough, R.S. Lloyd, Inhibition of DNA glycosylases via small molecule purine analogs, *PLoS One* 8 (2013) e81667.
- [17] S. Banerjee, K.L. Brown, M. Egli, M.P. Stone, Bypass of aflatoxin B₁ adducts by the *Sulfolobus solfataricus* DNA polymerase IV, *J. Am. Chem. Soc.* 133 (2011) 12556–12568.
- [18] P. Jaruga, E. Coskun, K. Kimbrough, A. Jacob, W.E. Johnson, M. Dizdaroglu, Biomarkers of oxidatively induced DNA damage in dreissenid mussels: a genotoxicity assessment tool for the Laurentian great lakes, *Environ. Toxicol.* 32 (2017) 2144–2153.
- [19] M. Dizdaroglu, E. Coskun, P. Jaruga, Measurement of oxidatively induced DNA damage and its repair, by mass spectrometric techniques, *Free Radic. Res.* 49 (2015) 525–548.
- [20] C. Zhu, L. Lu, J. Zhang, Z. Yue, J. Song, S. Zong, M. Liu, O. Stovicek, Y.Q. Gao, C. Yi, Tautomerization-dependent recognition and excision of oxidation damage in base-excision DNA repair, *Proc. Natl. Acad. Sci. U. S. A.* 113 (2016) 7792–7797.
- [21] E.F. Petterson, T.D. Goddard, C.C. Huang, G.S. Couch, D.M. Greenblatt, E.C. Meng, T.E. Ferrin, UCSF Chimera – a visualization system for exploratory research and analysis, *J. Comput. Chem.* 25 (2004) 1605–1612.
- [22] P. Emsley, B. Lohkamp, W.G. Scott, K. Cowtan, Features and development of coot, *Acta Crystallogr. D Biol. Crystallogr.* 66 (2010) 486–501.
- [23] D.A. Case, V. Babin, J.T. Berryman, R.M. Betz, Q. Cai, D.S. Cerutti, I.T.E. Cheatham, T.A. Darden, R.E. Duke, H. Gohlke, A.W. Goetz, S. Gusarov, N. Homeyer, P. Janowski, J. Kaus, I. Kolossváry, A. Kovalenko, T.S. Lee, S. LeGrand, T. Luchko, R. Luo, B. Madej, K.M. Merz, F. Paesani, D.R. Roe, A. Roitberg, C. Sagui, R. Salomon-Ferrer, G. Seabra, C.L. Simmerling, W. Smith, J. Swails, R.C. Walker,
- J. Wang, R.M. Wolf, X. Wu, P.A. Kollman, Amber14, University of California, San Francisco, 2014.
- [24] K.J. Karczewski, L.C. Francioli, G. Tiao, B.B. Cummings, J. Alfoldi, Q. Wang, R.L. Collins, K.M. Laricchia, A. Ganna, D.P. Birnbaum, L.D. Gauthier, H. Brand, M. Solomonson, N.A. Watts, D. Rhodes, M. Singer-Berk, E.G. Seaby, J.A. Kosmicki, R.K. Walters, K. Tashman, Y. Farjoun, E. Banks, T. Poterba, A. Wang, C. Seed, N. Whiffin, J.X. Chong, K.E. Samocha, E. Pierce-Hoffman, Z. Zappala, A.H. O'Donnell-Luria, E.V. Minikel, B. Weisburd, M. Lek, J.S. Ware, C. Vittal, I.M. Armean, L. Bergelson, K. Cibulskis, K.M. Connolly, M. Covarrubias, S. Donnelly, S. Ferriera, S. Gabriel, J. Gentry, N. Gupta, T. Jeandet, D. Kaplan, C. Llanwarne, R. Munshi, S. Novod, N. Petrillo, D. Roazen, V. Ruano-Rubio, A. Saltzman, M. Schleicher, J. Soto, K. Tibbetts, C. Tolonen, G. Wade, M.E. Talkowski, B.M. Neale, M.J. Daly, D.G. MacArthur, Variation across 141,456 human exomes and genomes reveals the spectrum of loss-of-function intolerance across human protein-coding genes, *bioRxiv* (2019) 531210.
- [25] A. Auton, L.D. Brooks, R.M. Durbin, E.P. Garrison, H.M. Kang, J.O. Korbel, J.L. Marchini, S. McCarthy, G.A. McVean, G.R. Abecasis, A global reference for human genetic variation, *Nature* 526 (2015) 68–74.
- [26] W. Zhang, H. He, M. Zang, Q. Wu, H. Zhao, L.L. Lu, P. Ma, H. Zheng, N. Wang, Y. Zhan, S. He, X. Chen, Z. Wu, X. Wang, J. Cai, Z. Liu, Z. Sun, Y.X. Zeng, C. Qu, Y. Jiao, Genetic features of aflatoxin-associated hepatocellular carcinoma, *Gastroenterology* 153 (249–262) (2017) e242.
- [27] J. Yeo, R.A. Goodman, N.T. Schirle, S.S. David, P.A. Beal, RNA editing changes the lesion specificity for the DNA repair enzyme NEIL1, *Proc. Natl. Acad. Sci. U. S. A.* 107 (2010) 20715–20719.
- [28] V. Bandaru, S. Sunkara, S.S. Wallace, J.P. Bond, A novel human DNA glycosylase that removes oxidative DNA damage and is homologous to *Escherichia coli* endonuclease VIII, *DNA Repair (Amst)* 1 (2002) 517–529.
- [29] I. Morland, V. Rolseth, L. Luna, T. Rognes, E. Seeberg, Human DNA glycosylases of the bacterial Fpg/MutM superfamily: an alternative pathway for the repair of 8-oxoguanine and other oxidation products in DNA, *Nucleic Acids Res.* 30 (2002) 4926–4936.
- [30] T.K. Hazra, T. Izumi, I. Boldogh, B. Imhoff, Y.W. Kow, P. Jaruga, M. Dizdaroglu, S. Mitra, Identification and characterization of a human DNA glycosylase for repair of modified bases in oxidatively damaged DNA, *Proc. Natl. Acad. Sci. U. S. A.* 99 (2002) 3523–3528.
- [31] K. Shinmura, H. Tao, M. Goto, H. Igarashi, T. Taniguchi, M. Maekawa, T. Takezaki, H. Sugimura, Inactivating mutations of the human base excision repair gene NEIL1 in gastric cancer, *Carcinogenesis* 25 (2004) 2311–2317.
- [32] S. Boiteux, E. Gajewski, J. Laval, M. Dizdaroglu, Substrate specificity of the *Escherichia coli* Fpg protein (formamidopyrimidine-DNA glycosylase): excision of purine lesions in DNA produced by ionizing radiation or photosensitization, *Biochemistry* 31 (1992) 106–110.
- [33] J. Tchou, V. Bodepudi, S. Shibutani, I. Antoshechkin, J. Miller, A.P. Grollman, F. Johnson, Substrate specificity of Fpg protein. Recognition and cleavage of oxidatively damaged DNA, *J. Biol. Chem.* 269 (1994) 15318–15324.
- [34] A. Karakaya, P. Jaruga, V.A. Bohr, A.P. Grollman, M. Dizdaroglu, Kinetics of excision of purine lesions from DNA by *Escherichia coli* Fpg protein, *Nucleic Acids Res.* 25 (1997) 474–479.
- [35] M. Dizdaroglu, C. Bauche, H. Rodriguez, J. Laval, Novel substrates of *Escherichia coli* Nth protein and its kinetics for excision of modified bases from DNA damaged by free radicals, *Biochemistry* 39 (2000) 5586–5592.
- [36] M. Dizdaroglu, J. Laval, S. Boiteux, Substrate specificity of the *Escherichia coli* endonuclease III: excision of thymine- and cytosine-derived lesions in DNA produced by radiation-generated free radicals, *Biochemistry* 32 (1993) 12105–12111.
- [37] A. Prakash, B.L. Carroll, J.B. Sweasy, S.S. Wallace, S. Double, Genome and cancer single nucleotide polymorphisms of the human NEIL1 DNA glycosylase: activity, structure, and the effect of editing, *DNA Repair (Amst)* 14 (2014) 17–26.
- [38] P.S. Pallan, L. Lei, C. Wang, M.R. Waterman, F.P. Guengerich, M. Egli, Research resource: correlating human cytochrome P450 21A2 crystal structure and phenotypes of mutations in congenital adrenal hyperplasia, *Mol. Endocrinol.* 29 (2015) 1375–1384.
- [39] H.H. Guo, J. Choe, L.A. Loeb, Protein tolerance to random amino acid change, *Proc. Natl. Acad. Sci. U. S. A.* 101 (2004) 9205–9210.
- [40] M. Forsbring, E.S. Vik, B. Dalhus, T.H. Karlsen, A. Bergquist, E. Schruppf, M. Bjoras, K.M. Boberg, I. Alseth, Catalytically impaired hMYH and NEIL1 mutant proteins identified in patients with primary sclerosing cholangitis and cholangiocarcinoma, *Carcinogenesis* 30 (2009) 1147–1154.